Notice to Readers: Inadvertent Intradermal Administration of Tetanus Toxoid--Containing Vaccines Instead of Tuberculosis Skin Tests

From MMWR Weekly July 30, 2004 / 53(29);662-664

Notice to Readers: Inadvertent Intradermal Administration of Tetanus Toxoid--Containing Vaccines Instead of Tuberculosis Skin TestsCDC and the Food and Drug Administration (FDA) have been notified about the potential for inadvertent administration of tetanus toxoid-containing vaccines (TTCVs) instead of tuberculin purified protein derivative (PPD) (Tubersol®, Aventis-Pasteur, Swiftwater, Pennsylvania; Aplisol®, Parkedale Pharmaceuticals, Rochester, Michigan) used for tuberculosis skin tests (TSTs). The Vaccine Adverse Event Reporting System (VAERS), a passive surveillance system jointly operated by CDC and FDA (1), detected clusters of medication errors in at least two states. These findings, along with another previously reported investigation involving the same error (2), suggest the need for health-care providers to take additional steps to minimize the risk for inadvertent intradermal injections of TTCVs.

In April 2004, five reports of medication error involving tetanus toxoid (TT) from a health-care provider were identified. Patients were vaccinated on three different dates; all experienced local reactions without complications. Another cluster reported to VAERS in June 2003 involved an undisclosed number of patients; a health-care provider confused tetanus and diphtheria toxoids (Td) vaccine for adult use (adsorbed) with PPD and administered Td intradermally. Patients with adverse reactions to these administrations had skin reactions interpreted as positive TSTs, which resulted in treatment with isoniazid (INH). Review of the lot numbers on products thought to be PPD revealed they were Td. Affected patients were identified and retested with PPD; all TSTs were negative. INH was discontinued, and no adverse reactions were observed.

As of March 2004, approximately 100 patients had been identified in reports of TTCV administration instead of PPD. A total of 21 states have reported both clusters and single cases. Vaccines substituted mistakenly for PPD include Td (n = 13 reports), TT (n = 12), and diphtheria and tetanus toxoids, (DT) adsorbed (n = five). For reports of Td, TT, and DT, products involved included those manufactured by Aventis-Pasteur and Wyeth (Collegeville, Pennsylvania) and vaccines from other unspecified manufacturers. CDC and FDA have initiated a full review of adverse events caused by inadvertent administration of vaccines and PPD products reported to VAERS and the FDA MedWatch Program. A preliminary review indicates that multiple vaccines other than TTCVs have been involved.

Similarities in packaging of PPD and TTCVs might have contributed to the medication errors (3,4). Both products require refrigeration and often are stored side by side. Lack of availability of Td in single-dose syringes, resulting in provider purchase of multiple-dose vials, was cited as a contributing factor to medication error in one cluster. Conversely, at least eight reports have been

documented of inadvertent substitution for vaccine products, resulting in intramuscular administration of PPD (FDA, unpublished data, 2004).

Health-care providers should consider ways to prevent vaccine misadministration. As more vaccines and combination products become available, the potential for medication errors might increase. Possible measures to prevent misadministration should include pharmacy dispensing of vaccines when feasible, physical separation of products, careful visual inspection and reading of labels, preparation of PPD for patient use only at time of testing, and improved record keeping of lot numbers of vaccines and other injectable products. Prevention of such errors through barcode scanning technology is the goal of a recent FDA rule requiring individual drug packages to have identifying barcodes (5). For health-care facilities that possess such technology, package scanning could help prevent errors made during pharmacy dispensing of products or during vaccine or PPD administration. In addition, the Product Identification Guide for Routine Vaccines is a helpful resource for distinguishing commonly used vaccine products; the guide can be ordered from the California Department of Health Services, telephone 619-594-5933. Adverse events associated with inadvertent vaccine administration can be reported to VAERS at http://www.vaers.org or by telephone, 800-822-7967. Adverse events after PPD administration can be reported to the FDA MedWatch program at http://www.fda.gov/medwatch or by telephone, 800-332-1088.

References

- 1. Chen RT, Rastogi SC, Mullen JR, et al. The Vaccine Adverse Event Reporting System (VAERS). Vaccine 1994;12:542--50.
- 2. Graham D, Dan B, Bertagnoll P, et al. Cutaneous inflammation caused by inadvertent intradermal administration of DTP instead of PPD. Am J Public Health 1981;71:1040--3.
- 3. Institute for Safe Medication Practices. Hazard alert! Confusion between tetanus diphtheria toxoid (Td) and tuberculin purified protein derivative (PPD) led to unnecessary treatment. Huntingdon Valley, Pennsylvania: Institute for Safe Medication Practices, 2003. Available at http://www.ismp.org/msaarticles/confusionprint.htm.
- 4. U.S. Food and Drug Administration. Mix up between Td and PPD. Rockville, Maryland: U.S. Department of Health and Human Services, U.S. Food and Drug Administration, 2003. Available at http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/psn/transcript.cfm
- 5. U.S. Food and Drug Administration. FDA rules requires bar codes on drugs and blood to help reduce errors. Rockville, Maryland: U.S. Department of Health and Human Services, U.S. Food and Drug Administration, 2004. Available at http://www.fda.gov/oc/initiatives/barcode-sadr.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S.

Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites. URL addresses listed in MMWR were current as of the date of publication.

Disclaimer All MMWR HTML versions of articles are electronic conversions from ASCII text into HTML. This conversion may have resulted in character translation or format errors in the HTML version. Users should not rely on this HTML document, but are referred to the electronic PDF version and/or the original MMWR paper copy for the official text, figures, and tables. An original paper copy of this issue can be obtained from the Superintendent of Documents, U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800. Contact GPO for current prices.